arXiv:hep-ph/0511044v1 4 Nov 2005

B DECAYS INTO $\pi\pi K$ AND $K\bar{K}K$: LONG DISTANCE AND FINAL-STATE EFFECTS

B. LOISEAU*

Laboratoire de Physique Nucléaire et de Hautes Énergies,[†] *Groupe Théorie,*
Univ. P. & M. Curie, 4 Pl. Jussieu, F-75252 Paris, France
E-mail: loiseau@lpnhep.in2p3.fr

A. FURMAN, R. KAMIŃSKI, L. LEŚNIAK

Department of Theoretical Physics,
The Henryk Niewodniczański Institute of Nuclear Physics,
Polish Academy of Sciences, 31-342 Kraków, Poland

The interplay of strong and weak decay amplitudes for $B \to \pi\pi K$ and $B \to K\bar{K}K$, with the $\pi\pi$ and $K\bar{K}$ pairs interacting in isospin-0 S-wave, is analyzed for $\pi\pi$ effective mass from threshold to 1.2 GeV. To improve agreement with experiment of a factorization approach with some QCD corrections, addition of long-distance contributions, called charming penguins is necessary.

1. Introduction

Hadronic two- and three-body B-meson weak decays are a rich source of information to study CP violation within the Standard Model (and beyond). On Dalitz plots, many resonances are distinguishable but with entangled interference patterns. Moreover the understanding of the final state interactions is weighty to attain accurate values of the Cabibbo-Kobayashi-Maskawa (CKM) matrix elements. We report here on our recent study[1] (FKLL) of $B \to \pi\pi K$ and $B \to K\bar{K}K$ decays. We consider the $\pi\pi$ or $K\bar{K}$ pairs, denoted by $(\pi\pi)_S$ or $(K\bar{K})_S$, to interact in isospin-0 S-wave in the $m_{\pi\pi}$ effective mass range from threshold to 1.2 GeV. We extend to the $f_0(980)$ case the approach of Gardner and Meißner[2] (GM) who have examined the effect of the $f_0(600)$ (or σ) on the $B^0 \to \pi^+\pi^-\pi^0$ decays. The decay amplitudes are calculated within the QCD factorization approach. Due to cancellations between penguin amplitudes, the $B \to f_0(980)K$ branching fractions are too small compared to data. We add long-distance contributions, called charming penguins. Expressions for the B decay amplitudes and description of the final state interactions, are given in section II. Comparison of our results to the data and some conclusions are presented in section III.

---

[*] Invited talk at the Third Asia-Pacific Conference on Few-Body Problems in Physics (APFB05), Nakhon Ratchassima, Thaïland, July 26-30, to be publised by the Scientific World Publishing Co.

[†] Unité de Recherche des Universités Paris 6 et Paris 7, associée au CNRS

2. B decay amplitudes and final state interactions

From the effective weak Hamiltonian, H_{eff}, and the operator product expansion, the B decay amplitude into mesons M_1 and M_2 is[3] (l stands for quarks d or s)

$$\langle M_1 M_2|H_{eff}|B\rangle = \frac{G_F}{\sqrt{2}} \sum_{q=u,c} V_{q,l}^{CKM} \left[\sum_{k=1}^{10} C_k(\mu) \langle M_1 M_2|O_k^{q,l}(\mu)|B\rangle \right]. \quad (1)$$

In (1) $G_F = 1.66 \times 10^{-5}$ GeV^{-2} is the Fermi constant, $V_{q,l}^{CKM}$ represents the CKM matrix-element factors and μ is a renormalization scale. The perturbative Wilson coefficients $C_k(\mu)$ characterize the short-distance physics. The non-perturbative hadronic matrix elements $\langle M_1 M_2|O_k^{q,l}(\mu)|B\rangle$ describe the long-distance physics. The local operators $O_k^{q,l}(\mu)$ *"effectively"* govern the decay. The main task of the theory is to compute these matrix elements in a reliable way. Beneke, Buchalla, Neubert and Sachrajda[4] have developed the following QCD factorization formula,

$$\langle M_1 M_2|O_k^{q,l}(\mu)|B\rangle \sim \langle M_1|J_1|0\rangle \langle M_2|J_2|B\rangle \left[1 + \sum_n r_n \alpha_s^n + O(\Lambda_{QCD}/m_b) \right]. \quad (2)$$

In (2) J_1 and J_2 are bilinear quark currents, $\langle M_1|J_1|0\rangle$ is the meson M_1 decay constant and $\langle M_2|J_2|B\rangle$ represents the transition form factor of the meson B into the meson M_2. The radiative corrections r_n have been calculated to order $n = 1$, α_s being the strong coupling constant. The term $O(\Lambda_{QCD}/m_b)$ in Eq. (2) represents power corrections and if set to zero, together with r_n, one recovers the naive factorization formula as applied by Bauer, Stech and Wirbel[5].

The possible quark line diagrams for the $B^- \to (\pi\pi)_S K^-$ and $B^- \to (K\bar{K})_S K^-$ are shown in Fig. 1 of FKLL[1]. For the B^0 decays there is no tree diagram (a), only penguin diagrams similar to (b) or (c) diagrams exist. Within the QCD factorization framework, just reminded above, the $B^- \to (\pi^+\pi^-)_S K^-$ decay amplitude is

$$\begin{aligned} \langle (\pi^+\pi^-)_S K^-|H_{eff}|B^-\rangle = \frac{G_F}{\sqrt{2}} \sqrt{\frac{2}{3}} \Big\{ & \chi \left[P(m_{\pi\pi})(U_a + U_b) + C(m_{\pi\pi}) \right] \Gamma_1^{n*}(m_{\pi\pi}) + \\ & + \left[Q(m_{\pi\pi}) U_c + \chi C(m_k) \right] \Gamma_1^{s*}(m_{\pi\pi}) \Big\}. \end{aligned} \quad (3)$$

The constant χ, to be fitted to reproduce the averaged branching fraction $\mathcal{B}[B^\pm \to f_0(980) K^\pm]$, is estimated to 30 GeV^{-1} from the $f_0(980)$ decay properties[1]. The non-strange and strange pion scalar form factors $\Gamma_1^n(m_{\pi\pi})$ and $\Gamma_1^s(m_{\pi\pi})$ describe the final-state interactions. In (3) $P(m_{\pi\pi})$ corresponds to the product of the kaon decay constant f_K times the transition form factor of B into $(\pi\pi)_S$, $F_0^{B\to(\pi\pi)_S}(M_K^2)$,

$$P(m_{\pi\pi}) = f_K (M_B^2 - m_{\pi\pi}^2) F_0^{B\to(\pi\pi)_S}(M_K^2). \quad (4)$$

The function $Q(m_{\pi\pi})$ is proportional to the product of the $(\pi\pi)_S$ "decay constant" times the transition form factor of B into K, $F_0^{B\to K}(m_{\pi\pi}^2)$, and reads[1,2],

$$Q(m_{\pi\pi}) = \frac{2\sqrt{2} B_0}{m_b - m_s} \left(M_B^2 - M_K^2 \right) F_0^{B\to K}\left(m_{\pi\pi}^2 \right), \quad (5)$$

with $B_0 = -\langle 0|q\bar{q}|0\rangle / f_\pi^2 = m_\pi^2/(m_u + m_d)$.

The functions U_a, U_b and U_c, given in terms of the coefficients[4,6,7,8] a_i of the decay operators, are calculated from the Wilson coefficients. They correspond to the contributions of Figs. 1(a), 1(b) and 1(c) of FKLL[1], respectively. One has $U_a = V_{ub}V_{us}^* a_1$, $U_b = V_{ub}V_{us}^* \left(a_4^u - a_6^u r\right) + V_{cb}V_{cs}^* \left(a_4^c - a_6^c r\right) = V_{ub}V_{us}^* \left[a_4^u - a_4^c + \left(a_6^c - a_6^u\right) r\right] + V_{tb}V_{ts}^* \left(a_6^c r - a_4^c\right)$ and $U_c = -V_{ub}V_{us}^* a_6^u - V_{cb}V_{cs}^* a_6^c = V_{ub}V_{us}^* \left(a_6^c - a_6^u\right) + V_{tb}V_{ts}^* a_6^c$. We have used the unitarity of the CKM matrix elements, $V_{ub}V_{us}^* + V_{cb}V_{cs}^* + V_{tb}V_{ts}^* = 0$. In the reduction of the four-quark operators, split into product of two-matrix elements (Eq. (2)), a Fierz transformation is needed for the penguin diagrams in order to match the flavor quantum numbers of the quark currents to those of the hadrons[6]. The chiral factor r is equal to $2M_K^2/[(m_b+m_u)(m_s+m_u)]$. Here and in Eq. (5), quark masses appear as one makes use of the Dirac equation. The a_1 and $a_{4,6}^{u,c}$ coefficients at next-to-leading order in α_s are given in Eq. (35) of Beneke and Neubert[7] (BN). In the numerical calculations we take the values of table III of de Groot, Cottingham and Whittingham[8] (GCW) at the scale $\mu = 2.1$ GeV. We do not include the small contributions from hard spectator interactions[7], annihilation and electroweak diagrams.

To improve their fit to charmless hadronic two-body B decays with pseudoscalar-pseudoscalar and pseudoscalar-vector decays, GCW[8], following Ciuchini et al.[9], introduce the charming penguin term $C(m)$, the parametrization of which is reminded in Eq. (6) of FKLL[1]. Expressions for the other $B \to (\pi\pi)_S K$ and for the $B \to (K\bar{K})_S K$ amplitudes are given in FKLL[1]. If one neglects the penguin contraction terms (see Eq. (35), (39) and (41) of BN[7]) then $a_{4,6}^u = a_{4,6}^c = a_{4,6}$ and our amplitude (3) is similar to that of the $B^- \to \sigma\pi^-$ decay[2] (see their Eq. (25)). Note the near cancellation[1] in the $b \to s$ transition of the two penguin contributions as $r \sim 1$ and $a_4^c \sim a_6^c$.

The $\Gamma_1^{n,s}(m)$ obey unitary $\pi\pi$ and $K\bar{K}$ coupled channel equations[1]. We use the solution A of the two-body scattering matrix of Kamiński, Leśniak and Loiseau[10]. Below the $K\bar{K}$ threshold $\Gamma_1^{n,s^*}(m) = R_1^{n,s}(m)\cos\delta_{\pi\pi}(m)e^{i\delta_{\pi\pi}(m)}$. $|\Gamma_1^{n,s}|$ is maximum for $\delta_{\pi\pi} \simeq 180°$, which corresponds to the $f_0(980)$ production. The $R_1^{n,s}(m)$, depicting the formation of mesons prior to rescattering, have been obtained by Meißner and Oller[11].

3. Results and conclusions

Our results with the charming-penguin $C(m)$ terms of GCW[8], model I, and of Ciuchini, Franco, Martinelli, Masiero, Pierini and Silvestrini[12] (CFMMPS), model II, are compared to the data[13,14,15,16] in Table 1. Values of all parameters used are given in FKLL[1]. The effective $m_{\pi\pi}$ mass distribution for $B^\pm \to \pi^+\pi^- K^\pm$ of our model II is compared with Belle[13] in Fig. 1. The sharp maximum near 1 GeV is from the $f_0(980)$. Near 0.5 GeV the broad bump is related to the $f_0(600)$. Events near 0.8 GeV are due to the $B^\pm \to \rho^0 K^\pm$ decay not considered in our model. Similar agreement with $m_{\pi\pi}$ effective mass distributions of BaBar and Belle



Figure 1. Comparison of our model II with Belle data[13]

Table 1. Average branching fractions $\mathcal{B}$ in units of 10^{-6}, direct asymmetries $\mathcal{A}_{CP}$, $\mathcal{A}$ and $\mathcal{S}$ parameters of our model compared to data. Model errors come from $C(m)$ parameters fitted by GCW[8], model I, and CFMMPS[12], model II. For a qualitative comparison we give in parenthesis the experimental $B \to K\bar{K}K$ decay observables obtained by considering the full range of the $K\bar{K}$ invariant mass rather than the upper limit[1] of 1.1 GeV. For data[14,16] we only quote statistical errors.

B decay mode		Average HFAG's values[15]	Model I $\chi = 33.5$ GeV^{-1}	Model II $\chi = 23.5$ GeV^{-1}
$B^{\pm} \to f_0(980)K^{\pm}$,	$\mathcal{B}$	$8.49^{+1.35}_{-1.26}$	8.49 (fit)	8.46 (fit)
$f_0 \to \pi^+\pi^-$	$\mathcal{A}_{CP}$	-0.02 ± 0.07	-0.52 ± 0.12	0.20 ± 0.20
$B^0 \to f_0(980)K^0$,	$\mathcal{B}$	6.0 ± 1.6	5.9 ± 1.6	5.8 ± 2.8
$f_0 \to \pi^+\pi^-$	$\mathcal{A}$	-0.14 ± 0.22	0.01 ± 0.10	0.0004 ± 0.0010
	$\mathcal{S}$	-0.39 ± 0.26	-0.63 ± 0.09	-0.77 ± 0.0004
$B^{\pm} \to \left(K^+K^-\right)_S K^{\pm}$	$\mathcal{B}$	$< 2.9^{13}$	1.8 ± 0.4	1.7 ± 0.7
	$\mathcal{A}_{CP}$	no data	-0.44 ± 0.12	0.29 ± 0.21
$B^0 \to \left(K^+K^-\right)_S K^0_S$	$\mathcal{B}$	no data	1.1 ± 0.3	1.2 ± 0.5
	$\mathcal{A}$	(-0.09 ± 0.10)	0.01 ± 0.10	0.001 ± 0.001
	$\mathcal{S}$	$(-0.55 \pm 0.22)^{14}$ $(-0.74 \pm 0.27)^{16}$	-0.64 ± 0.09	-0.77 ± 0.0006
$B^0 \to \left(K^0_S K^0_S\right)_S K^0_S$	$\mathcal{B}$	no data	1.1 ± 0.3	1.2 ± 0.5
	$\mathcal{A}$	(0.41 ± 0.21)	0.01 ± 0.10	0.001 ± 0.001
	$\mathcal{S}$	(-0.26 ± 0.34)	-0.64 ± 0.09	-0.77 ± 0.0006

for charged and neutral B decays can be seen in the figures shown in FKLL[1].

To conclude, our effective mass distributions and branching fractions compare well with data. The long distance contribution $C(m)$ is important. If $C(m) = 0$,

$\mathcal{B}(B^0 \to \pi^+\pi^- K^0_S)$ is too small by a factor ~ 18 and $\mathcal{B}(B^\pm \to \pi^+\pi^- K^\pm)$ by a factor ~ 4. If charming penguins are included the agreement with data is good with a χ compatible with the $f_0(980)$ properties. The recent determination[15] of $\mathcal{A}_{CP} = -0.02 \pm 0.07$ for $B^\pm \to f_0(980)K^\pm,\ f_0 \to \pi^+\pi^-$, should help to determine better the charming-penguin $C(m)$ parameters. In progress we are adding the P-wave contribution which will allow us to describe the ρ-meson region.

Acknowledgments

We acknowledge good advices from B. El-Bennich. This work benefits from IN2P3-Polish Laboratories Convention (project N^0 99-97).

References

1. A. Furman, R. Kamiński, L. Leśniak and B. Loiseau, Phys. Lett. **B622**, 207 (2005), *Long-distance effects and final state interactions in* $B \to \pi\pi$ *and* $B \to K\bar{K}K$ *decays.*
2. S. Gardner and U.-G. Meißner, Phys. Rev. **D65**, 094004 (2002), *Rescattering and chiral dynamics in* $B \to \rho\pi$ *decay.*
3. *The BABAR Physics Book: Physics at an Asymmetric B Factory* by BABAR collaboration, P. F. Harrison and H. R. Quinn, Editors, report SLAC-R-504, Oct. 1998 (see in particular chapters 2 and 10).
4. M. Beneke, G. Buchalla, M. Neubert and C. T. Sachrajda, Phys. Rev. Lett. **83**, 1914 (1999), *QCD factorization for* $B \to \pi\pi$ *decays: strong phases and* CP *violation in the heavy quark limit;*
5. M. Bauer, B. Stech and M. Wirbel, Z. Phys. **C34**, 103 (1987), *Exclusive non-leptonic decays of* $D-$, D_s- *and* $B-$ *mesons.*
6. A. Ali, G. Kramer and C.-D. Lü, Phys. Rev. **D58**, 094009 (1998), *Experimental tests of factorization in charmless non leptonic two-body* B *decays.*
7. M. Beneke, M. Neubert, Nucl. Phys. **B675**, 333 (2003), *QCD factorization for* $B \to PP$ *and* $B \to PV$ *decays.*
8. N. de Groot, W. N. Cottingham and I. B. Whittingham, Phys. Rev. **D68**, 113005 (2003), *Factorization fits and the unitarity triangle in charmless two-body* B *decays.*
9. M. Ciuchini, E. Franco, G. Martinelli and L. Silvestrini, Nucl. Phys. **B501**, 271 (1997), *Charming penguins in* B *decays.*
10. R. Kamiński, L. Leśniak and B. Loiseau, Phys. Lett. **B413**, 130 (1997), *Three channel model of meson-meson scattering and scalar meson spectroscopy.*
11. U.-G. Meißner and J . A. Oller, Nucl. Phys. **A679**, 671 (2001), $J/\psi \to \phi\pi\pi(K\bar{K})$ *decays, chiral dynamics and OZI violation.*
12. M. Ciuchini, E. Franco, G. Martinelli, A. Masiero, M. Pierini and L. Silvestrini, talk at Rencontre de Moriond, March 2004, hep-ph/0407073, *Two-Body noleptonic B decays in the Standard Model and beyond.*
13. A. Garmash *et al.*, Belle Collaboration, Phys. Rev. **D71**, 092003 (2005), *Dalitz analysis of the three-body charmless decays* $B^+ \to K^+\pi^+\pi^-$ *and* $B^+ \to K^+K^+K^-$.
14. B. Aubert *et al.,* BaBar Collaboration, Phys. Rev. **D71**, 091102(R) (2005), *Measurement of CP asymmetries in* $B^0 \to \phi K^0$ *and* $B^0 \to K^+K^-K^0_S$ *decays.*
15. J. Alexander *et al.*, The Heavy Flavor Averaging Group (HFAG), http://www.slac.stanford.edu/xorg/hfag/index.html.
16. K.-F. Chen *et al.*, Belle Collaboration, Phys. Rev. **D72**, 012004 (2005), *Time-dependent* CP*-violating asymmetries in* $b \to s\bar{q}q$ *transitions.*